

AIM® EC HERBICIDE

DORMANT AND/OR BETWEEN CUTTINGS

PRODUCT ATTRIBUTES	BENEFITS
Fast Activity	Aim® EC herbicide works fast. Whether used alone or in a tank mix with other labeled herbicides, Aim EC herbicide will burn down labeled weeds in just a few days. Enhances glyphosate herbicide activity in Roundup Ready® alfalfa fields and helps manage shifting weed populations.
Increased Weed Spectrum	Excellent activity on velvetleaf, morningglory species, lambsquarters, nightshade, pigweed and waterhemp. Improves control of kochia, Russian thistle and field bindweed.
PPO Chemistry	The mode of action of Aim EC herbicide results in thorough desiccation to improve weed control and alleviate crop competition.
Application Flexibility	With its wide window of application, Aim EC herbicide can be used as a dormant spray or between cuttings to maximize weed control. Aim EC herbicide can be tank mixed with many herbicides for improved weed spectrum.

APPLICATION RECOMMENDATIONS	
Use Rates:	0.5 – 2.5 fl. oz./A
Active Ingredient	Carfentrazone-ethyl
Tank-Mix Recommendations:	<ul style="list-style-type: none"> May be applied to dormant alfalfa or alfalfa stubble alone or in combination with other registered herbicides. Residual herbicides may also be used according to their labels. Do not drift spray mixture on desirable vegetation. Target weeds of 4 inches or less for optimum control. Additional surfactant is required. MSO at 1% v/v is preferred. Use a minimum of 10 gallons finished spray per acre by ground and 5 gallons by air. Maximum seasonal use rate is 2.5 fl. oz./A of Aim EC herbicide. PHI: 21 days for forage or hay. After application of this product, you may rotate to any Aim EC herbicide labeled crop at anytime.

ADJUVANT GUIDELINES	
<ul style="list-style-type: none"> COC or an MSO at 1% v/v is recommended under dry conditions, heavy weed pressure. Add AMS at 2-4 lbs./A or UAN at 2-4% v/v is allowed. 	
WEEDS CONTROLLED	
0.5 – 1.0 fl. oz./A	1.5 – 2.5 fl. oz./A
<ul style="list-style-type: none"> Bedstraw, catchweed Cheeseweed Flixweed Lambsquarters (up to 3") Fiddleneck, coast Mallow, common Mustard, tansy Nightshade, black Pennycress, field Pigweed, redroot Rocket, London Velvetleaf Wallflower, bushy 	<ul style="list-style-type: none"> Amaranthus, spp. Bittercress Buckwheat, wild Filaree, redstem Kochia Lambsquarters Mustard, tumble Meadowfoam Mustard, tansy Nightshade, hairy Shepherd's purse Sowthistle, annual Speedwell, ivyleaf Spurry, corn

WEED CONTROL 60 DAT – DORMANT APPLICATION – 2013 | BOYDSTON, WASHINGTON STATE UNIVERSITY

